



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|-----------|---|
| <p>(51) International Patent Classification ⁶ : C07D 213/00, 213/74, 231/1285, 239/91, 285/28, 295/14, 307/52, 311/00, 311/22, 333/22, 409/12, 417/12, A61K 31/405, 31/245, 31/63</p> | <p>A2</p> | <p>(11) International Publication Number: WO 98/09948</p> <p>(43) International Publication Date: 12 March 1998 (12.03.98)</p> |
| <p>(21) International Application Number: PCT/EP97/04774</p> <p>(22) International Filing Date: 2 September 1997 (02.09.97)</p> <p>(30) Priority Data: MI96A001821 4 September 1996 (04.09.96) IT</p> <p>(71) Applicant (for all designated States except US): NICOX S.A. [FR/FR]; 45, avenue Kléber, F-75116 Paris (FR).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): DEL SOLDATO, Piero [IT/IT]; Via Toti, 22, I-20052 Monza (IT). SANNICOLO', Francesco [IT/IT]; Alzaia Naviglio Grande, 46, I-20148 Milano (IT).</p> <p>(74) Agents: SAMA, Daniele et al.; Sama Patents, Via G.B. Morgani 2, I-20129 Milano (IT).</p> | | <p>(81) Designated States: AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p> |
| <p>(54) Title: NITRIC ESTER DERIVATIVES AND THEIR USE IN URINARY INCONTINENCE AND OTHER DISEASES</p> <p>(57) Abstract</p> <p>Use of the following groups of compounds or their compositions for the preparation of medicaments for the treatment of urinary incontinence, such compounds having general formula: A-X₁-NO₂ or their salts, where A = R(COX)_t and where t is an integer 0 or 1; X = O, NH, NR_{1C}, where R_{1C} is a linear or branched alkyl having from 1 to 10 C atoms; R is (IA) where t = 1 and X₁ is equal to -YO- where Y is a C₁-C₂₀ alkylene, C₅-C₇ cycloalkyl or oxyalkyl derivatives.</p> <div data-bbox="1036 1150 1421 1480" style="text-align: center;"> <p>(IA)</p> </div> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

NITRIC ESTER DERIVATIVES AND THEIR USE IN URINARY INCONTINENCE AND OTHER DISEASES

The present invention relates to new medicaments to be used in urinary disorders. These disorders are generally grouped in one single functional pathology class and are characterized by several symptoms, including changes in micturition (like for example incontinence), changes in urinary output (like for example polyuria, oliguria, anuria), changes in the appearance of urine (like for example hematuria), edema (like for example anasarca), pain (like for example bladder pain).

The invention relates to new compounds having superior efficacy in the treatment of some forms of urinary incontinence (anti-incontinence compounds) or edema (diuretics) and which appear to be well tolerated by the body.

In particular, it is known that urinary incontinence can be considered a disorder of micturition control resulting from a lesion or dysfunction of the lower urinary tract. In particular, the urinary bladder smooth muscle called detrusor muscle and the internal (smooth muscle) and external (striated muscle) urethral sphincters are involved. See for example Ferguson D. and Christopher N., *Urinary bladder function and drug development*, Trends in Pharmacological Sciences, 1996, 17, 161-165. This publication reports

that there are various types of incontinence characterised by different causes and symptoms. In particular, the following can be mentioned:

- stress incontinence, which is the discharge of small amounts of urine due to increased intraabdominal pressure caused, for example, by cough or an effort. Stress incontinence is due to a change in vesicourethral angle and relaxation of the urethral sphincter muscle. Stress incontinence is frequent in women, particular multipara women.
- urge incontinence, which is the inability to control the urinary bladder and manifests itself with a sudden and urgent stimulus to urinate. Urge incontinence is due to intermittent contraction of the urinary bladder muscle for no apparent cause (detrusor instability) or caused by interstitial cystitis or other inflammatory phenomena which lead to urinary bladder hyperexcitability. It seems that changes in urinary bladder innervation are present in all these cases;
- incontinence from urinary bladder overdistention, which occurs in case of chronic urinary retention due to obstructive causes. The urinary bladder never empties completely, resulting in continous discharge of small

amounts of urine.

- total incontinence, which is a complete lack of urinary bladder control due to inability of controlling the sphincters. It is the result of severe neurologic damages.

In the known art, the available therapies are based on three different approaches, see for example the above publication and Anderson K.E., *Pharmacology of lower urinary tract smooth muscles and penile erectile tissues*, *Pharmacological Reviews*, 1993, 45, 253-308:

- reduced detrusor activity,
- changed sensory nervous transmission,
- changed urethral resistance.

According to the first approach, detrusor contraction is stimulated by the parasympathetic system and acetylcholine is the main mediator. Therefore, anticholinergic agents are used to reduce vesical hyperactivity. However, these are effective but of limited use due to the systemic anticholinergic side effects including for example dry mouth, constipation and tachycardia. Taking into account that vesical irritability is often associated with urinary bladder obstructive disease, the administration of anticholinergic agents risks triggering an acute urinary retention crisis.

Another pharmacological approach to reduce detrusor activity includes the use of medicaments which help opening potassium channels or calcium antagonists which relax the smooth muscle. However, there are disadvantages such as a marked hypotensive action due to a nonspecific vasodilator effect produced by these agents.

An additional pharmacological measure to reduce detrusor activity consists of the use of prostaglandin synthesis inhibitors which were tested in some detrusor hyperactivity and enuresis cases with promising results but giving major side effects. Their use is based on the fact that numerous prostaglandins were found to be synthesised in the urinary bladder following nervous stimulation and some of them seem to act as mediators of detrusor contraction. Additionally, some prostaglandines may be involved in severe urge incontinence and vesical hyperactivity events during some inflammatory disease of the urinary tract.

Therefore, nonsteroidal anti-inflammatory drugs are potentially useful in lowering the urinary bladder excitability threshold, and are thus effective in cases of detrusor instability. Unfortunately, they have the disadvantage of being little tolerated at active doses, especially in the gastrointestinal tract.

Likewise, NO synthetase enzyme inhibitors can prevent hyperexcitability of the urinary bladder and hyperalgesia resulting from inflammatory events such as interstitial cystitis; see Rice A.S.C., *Topical spinal administration of a nitric oxide synthase inhibitor prevents the hyperreflexia associated with a rat model of persistent visceral pain*, Neuroscience Letters, 1995, 187, 111-114. However, there are currently no agents of this type which can be used therapeutically due to a relative nonspecificity of their pharmacological profile.

The second approach, which consists of changing sensory nervous transmission (whenever urinary incontinence results from lesions of the nervous system), includes the use of drugs which act on neurotransmission, for example gamma-aminobutyric acid (GABA), or peptides, or purines, which are important neurotransmitters in the urinary tract.

Studies are also known which use capsaicin for intravesical instillation with sometimes satisfactory results. However, this treatment has limited clinical applications due to its transient effect, which, in addition, can be obtained only by local use.

The third approach considers the fact that muscle tone in the urethra is mediated by different neurotransmission


systems, including for example the adrenergic system, by stimulation of α -receptors, α -agonist medicaments, which increase the pressure borne by the urethra, are used to change urethral resistance sometimes with satisfactory results. However, the use of these compounds involves some risks, as in the case of urinary bladder obstructive disease where even alpha-antagonists are used. In these cases a sphincter hyperactivity is observed, which prevents regular urinary bladder voiding causing urge incontinence. Also in this case, as in the first approach described above, severe side effects of a hypotensive type related to the α -antagonistic activity in the cardiocirculatory system are observed.

To increase urethral resistance in women with stress incontinence, an oestrogen based therapy which was found to be efficacious in increasing intraurethral pressure and in changing the structure of mucous membrane, vessel and connective, is used. Good results were observed combining treatment with α -agonists with oestrogen treatment. However, the well known side effects which occur when oestrogen treatment is used must be reported.

So far, commercial pharmaceutical preparations resolve the problem only in a limited number of cases. However, they generally cause side effects, even somewhat severe.

The Applicant has unexpectedly and surprisingly found that the particular classes of compounds described below can be beneficially used in the treatment of the various types of urinary incontinence described above, as they exhibit a pharmacological profile superior to that of the known preparations used for this type of disease.

An object of the present invention is the use for the treatment of urinary incontinence of the following classes of compounds, having the general formula:




or their salts, wherein

A = R(COX)_t where t is an integer 0 or 1;


X = O, NH, NR_{1C} where R_{1C} is a linear or branched alkyl having from 1 to 10 C atoms;

R is chosen from the following groups:

* Group IA), wherein t = 1



(IAa)



wherein

R_{II5} is H, a linear or whenever possible branched C_1 - C_3 , alkyl;

R_{II6} has the same meaning as R_{II5} , or when R_{II5} is H it can be benzyl;

R_{II1} , R_{II2} and R_{II3} , equal or different, are hydrogen, linear or whenever possible branched C_1 - C_6 alkyl or C_1 - C_6 alkoxy, or Cl, F, Br;

R_{II4} is R_{II1} or bromine;


preferred are the compounds where R_{II1} , R_{II2} and R_{II4} are H, and R_{II3} is Cl, and R_{II3} is in the ortho position to NH; R_{II5} and R_{II6} are H, X is equal to O, and X_1 is $(CH_2-CH_2-O)_2$;

(IAb) is the residue of 2-[[2-methyl-3-(trifluoromethyl)phenyl]amino]-3-pyridinecarboxylic acid and when -COOH is present it is known as flunixin.

The compounds preferred are those where $X = O$;

* IIA) chosen from the following

wherein when $t = 1$,




wherein


R_{2a} and R_{3a} are H, a linear or whenever possible branched substituted or non-substituted C_1 - C_{12} alkyl, allyl, with the proviso that when one of the two is allyl the other is H; preferably R_{2a} is H, alkyl has from 1 to 4 C atoms, R_{3a} is H;

R_{1a} is chosen from


II Aa)




(II)




(XXI)




(IV)




(VII)




(XXV)




(VI)




(VIII)



(IX)



(X)



(III)

wherein :

- in the compounds of formula (IV), residue of ketoprofen:

R_{III1} is H, SR_{III3} where R_{III3} contains from 1 to 4 C atoms linear or whenever possible branched;

R_{III2} is H, hydroxy;

preferred are the compounds where R_{III1} and R_{III2} are H, R_{3a} is H, and R_{2a} is methyl, $X = O$;

- in the compounds of formula (XXI), residue of carprofen:

R_{xxio} is H, a linear or whenever possible branched alkyl having from 1 to 6 carbon atoms, a C_1-C_6 alkoxycarbonyl bound to a C_1-C_6 alkyl, a C_1-C_6 carboxyalkyl, a C_1-C_6 alkanoyl, optionally substituted with halogen, benzyl or halobenzyl, benzoyl or halobenzoyl;

R_{xxi} is H, halogen, hydroxy, CN, a C_1-C_6 alkyl optionally containing OH groups, a C_1-C_6 alkoxy, acetyl, benzyloxy, SR_{xxi2} where R_{xxi2} is C_1-C_6 alkyl; a perfluoroalkyl having from 1 to 3 C atoms, a C_1-C_6 carboxyalkyl optionally containing OH, NO_2 , sulphamoyl, dialkyl sulphamoyl groups with the alkyl having from 1 to 6 C atoms, or difluoroalkylsulphonyl with the alkyl having from 1 to

3 C atoms;

R_{xxi1} is halogen, CN, a C_1 - C_6 alkyl containing one or more OH, a C_1 - C_6 alkoxy, acetyl, acetamide, benzyloxy groups, SR_{III3} is as above defined, a perfluoroalkyl having from 1 to 3 C atoms, hydroxy, a carboxyalkyl having from 1 to 6 C atoms, NO_2 , amino, a mono- or dialkylamino having from 1 to 6 C atoms, sulphamoyl, a dialkyl sulphamoyl having from 1 to 6 C atoms, or difluoroalkylsulphamoyl as above defined; or R_{xxi} together with R_{xxi1} is an alkylene dioxy having from 1 to 6 C atoms;

preferred are the compounds where R_{xxio} is H, the connection bridge is at position 2, R_{xxi} is H, R_{xxi1} is chlorine and is in para position to nitrogen;

R_{3a} is H, R_{2a} is methyl and X is O;

- in the compounds of formula (XXXV), residue of thiaprofenic acid:


Ar is phenyl, hydroxyphenyl optionally mono or polysubstituted with halogen, an alkanoyl or alkoxy having from 1 to 6 C atoms, a trialkyl having from 1 to 6 C atoms, preferably from 1 to 3 C atoms, cyclopentyl o-hexyl o-heptyl, heteroaryl, preferably thienyl, furyl optionally containing OH, pyridyl;

the preferred compounds of formula (XXXV) are those


- where Ar is phenyl, R_{3a} is H, R_{2a} is methyl and X is O;
- in the compounds of formula (II), residue of suprofen, of which the preferred, where $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$, has been shown, its equivalents, as described and obtained in patent USP 4,035,376, herein incorporated by reference, can be used.
 - in the compounds of formula (VI), of which the preferred, indoprofen, when R_{2a} is CH_3 or indobufen, when R_{2a} is equal to H and $R_{3a} = CH_3$ and $X = O$, have been shown, its equivalents, as described and obtained in accordance with patent USP 3,997,669, herein incorporated by reference, can be used.
 - in the compounds of formula (VIII), of which the preferred etodolac, when $R_{3a} = R_{2a} = H$ and $X = O$, has been shown, its equivalents as described and obtained according to patent USP 3,843,681, herein incorporated by reference, can be used;
 - in the compounds of formula (VII), of which the preferred, fenoprofen, when $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$, has been shown, its equivalents, as described and obtained according to patent USP 3,600,437, herein incorporated by reference, can be used;

- in the compounds of formula (III),
of which the preferred, fenbufen, when $R_{3a} = R_{2a} = H$ and $X = O$, has been shown, its equivalents, as described and obtained according to patent USP 3,784,701, herein incorporated by reference, can be used;
- in the compounds of formula (X), residue of tolmetin, when $R_{3a} = R_{2a} = H$ and $X = O$, its equivalents, as described and obtained according to patent FR 1.574.570, herein incorporated by reference, can be used;
- in the compounds of formula (IX), residue of flurbiprofen, when $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$, its equivalents as described in the known art can be used;


II Ab)




IIIa)




(XXX)




(XXXI)




(XXXII)



(XXXIII)



(XXXVI)




where the meanings are as follows:

- when IIIa) contains $-\text{CH}(\text{CH}_3)-\text{COOH}$ it is known as prano-profen: α -methyl-5H-[1] benzopyran [2,3-b]pyridine-7-acetic acid. In the preferred compounds $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$ and $\text{X} = \text{O}$;
- when residue (XXX) contains $-\text{CH}(\text{CH}_3)-\text{COOH}$ it is known as bermoprofen: dibenz [b,f] oxepin-2-acetic acid. The preferred compound is that with $\text{X} = \text{O}$, $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$;
- residue (XXXI) is known as CS-670, 2-[4(2-oxo-1-cyclohexylidenemethyl)phenyl]propionic acid, when the radical is $-\text{CH}(\text{CH}_3)-\text{COOH}$. The preferred compound has $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}$ and $\text{X} = \text{O}$;
- residue (XXXII) derives from the known pemedolac which contains group $-\text{CH}_2\text{COOH}$. The compound preferred has $\text{R}_{2a} = \text{R}_{3a} = \text{H}$ and $\text{X} = \text{O}$;
- when residue (XXXIII) is saturated with $-\text{CH}_2\text{COOH}$ it is

known as pyrazolac: 4-(4-chlorophenyl)-1-(4-fluorophenyl)3-pyrazolyl acid derivatives. The preferred compounds have $R_{2a} = R_{3a} = H$ and $X = O$;

- when residue (XXXVI) is saturated with $-CH(CH_3)-COO-$ it is known as zaltoprofen. When the residue is saturated with a hydroxy or amine group or the acid salts, the compounds are known as dibenzothiepin derivatives. The compounds preferred have $R_{2a} = H$, $R_{3a} = CH_3$ and $X = O$;
- when residue (XXXVII) is CH_2-COOH it derives from the known mofezolac: 3,4-di(p-methoxyphenyl)isoxazol-5-acetic acid. The preferred compounds include $R_{2a} = R_{3a} = H$, $t = 1$, $X = O$.


* Group III A), where $t = 1$,




where

R_{IVd} and R_{IVd1} are at least one H and the other a linear or whenever possible branched C_1-C_6 alkyl, preferably C_1 and C_2 , or difluoroalkyl with the alkyl having from 1 to 6 C atoms, preferred is C_1 , or R_{IVd} and R_{IVd1} jointly form a methylene group;


R_{IV} has the following meaning:



(II)



(X)



(III)

where the compounds of group III A) have the following meanings:

- in the compounds of formula (II):

R_{IV-II} is an alkyl having from 1 to 6 C atoms, a cycloalkyl having from 3 to 7 C atoms, an alcoxymethyl having from 1 to 7 C atoms, a trifluoroalkyl having from 1 to 3 C atoms, vinyl, ethynyl, halogen, an alkoxy having from 1 to 6 C atoms, a difluoroalkoxy with the


alkyl having from 1 to 7 C atoms, an alkoxymethyloxy having from 1 to 7 C atoms, an alkylthiomethyloxy with the alkyl having from 1 to 7 C atoms, an alkylmethylthio with the alkyl having from 1 to 7 C atoms, cyano, difluoromethylthio, a substituted phenyl- or phenylalkyl with the alkyl having from 1 to 8 C atoms; preferably R_{IV-II} is CH_3O , R_{IVd} is H and R_{IVd1} is CH_3 , and is known as the residue of naproxen;

X = NH and X_1 is equal to $(CH_2)_4$ or $(CH_2CH_2O)_2$; also preferred is the same compound where X is equal to O;

- in the compounds of formula (X), for which the residue of loxoprofen has been shown, the residues described in USP 4,161,538 herein incorporated by reference can be used as equivalents; preferred are the compounds where R_{IVd} is H and R_{IVd1} is CH_3 , X = NH and X_1 is equal to $(CH_2)_4$ or $(CH_2CH_2O)_2$; also preferred is the same compound where X is equal to O;


- in the compounds of formula (III):

R_{IV-III} is a C_2 - C_5 alkyl, even branched whenever possible, a C_2 and a C_3 alkyloxy, allyloxy, phenoxy, phenylthio, a cycloalkyl having from 5 to 7 C atoms, optionally substituted at position 1 by C_1 - C_2 alkyl; preferred is the compound where R_{IV-III} is



and $R_{IVd} = \text{H}$, R_{IVd1} is CH_3 , a compound known as the residue of ibuprofen; $X = \text{NH}$ and X_1 is equal to $(\text{CH}_2)_4$ or $(\text{CH}_2\text{CH}_2\text{O})_2$; also preferred is the same compound where X is equal to O ;

* Group IV A)



(IV)


where $A = \text{RCOO}$, $t = 1$,

of which the residue of the known indomethacin has been shown, its equivalents as described and obtained in patent USP 3,161,654 herein incorporated by reference can be used;


* Group V A) chosen from the following:

- V Aa) fenamates chosen from the following,


where $t = 1$




(V Aa1)




(V Aa2)



(V Aa3)




(V Aa4)




(V Aa5)

-V Ab), derivatives of niflumic acid, where $t = 1$:




(V Ab1)


- V Ac), COX₂ inhibitors, where $t = 0$ and R is as follows:




(V Ac1)




(V Ac2)






(V Ac3)




(V Ac4)




- V Ad) Derivatives of diuretics when $t = 1$ and R is as follows:






(V Ad4)


- V Ae) Derivatives of diuretics when $t = 0$ and R is as follows:




(V Ae1)




(V Ae2)




(V Ae3)




(V Ae4)




(V Ae5)




(V Ae6)




(V Ae7)




(V Ae8)




(V Ae9)



(V Ae10)



(V Ae11)



(V Ae12)

where the meaning in group V A) is as follows:

in V Aa):

- in compounds (V Aa1) the residue of enfenamic acid, 2-[(2-phenylethyl)amino]benzoic acid, where COOH was substituted according to the present invention, has been shown.

This can be prepared according to the Indian patents 103.066 and 114.805, herein incorporated by reference. Equivalent products containing various substituents as described in said patents can be used, too.

- In compounds (V Aa2) the residue of flufenamic acid, 2-[[3-(trifluoromethyl)phenyl]-amino]benzoic acid, where COOH was substituted according to the present invention, has been shown.

This can be prepared according to Wilkinson's article, Finar, J.Chem.Soc. 1948, 32, herein incorporated by reference. Any equivalent product containing various substituents as described in said article can be used, too.

- In compounds (V Aa3) the residue of meclofenamic acid, 2-[(2,6-dichloro-3-methylphenyl)amino]benzoic acid where COOH was substituted according to the present invention, has been shown.

This can be prepared according to the German patent DE 1.149.015 and USP 3.313.848 herein incorporated by reference. Any equivalent product containing various substituents as described in said patents can be used, too.

- In compounds (V Aa4) the residue of mefenamic acid, 2-[(2,3-dimethylphenyl)amino]benzoic acid where COOH was substituted according to the present invention, has been shown.

This can be prepared according to the Belgian patent 605.302, herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Aa5) the residue of tolfenamic acid, 2-[(3-chloro-2-methylphenyl)amino]benzoic acid where COOH was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3.313.848, herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

In V Ab):

- in compounds (V Ab1) the residue of niflumic acid,

2-[[3-(trifluoromethyl)phenyl]amino]-3-pyridine carboxylic acid, where COOH was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3.415.834, herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

In V Ac):

- in compound (V Ac1) R_{vac1} attached to the oxygen atom in position 2 of the benzene ring of N-(4-nitrophenyl)methansulphonamide can be phenyl or cyclohexane. When R_{vac1} is phenyl the residue is that of nimesulide.

This can be prepared according to patent USP 3,840,597 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ac2) the residue of 3-formylamino-7-methylsulfonylamino-6-phenoxy-4H-1-benzopyran-4-one was substituted according to the present invention, has been shown.

This can be prepared according to patent DE 3834204 herein incorporated by reference. Any equivalent product containing various substituents as described in

said patent can be used, too.

- In compounds (V Ac3) the atom X₄ that links the radical 2,4-difluorothiophenyl to position 6 of the indanone ring of the residue 5-methanesulfonamido-1-indanone can be sulfur or oxygen.

This can be prepared according to WO 9413635 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

In compounds (V Ac4) the residue of celecoxib 4-[5(4-methylphenyl)-3-(trifluoromethyl)pyrazol-1-yl] benzenesulphonamide was substituted according to the present invention, has been shown.

This can be prepared according to patent WO 9427980 herein incorporated by reference. Any equivalent product containing various substituents e.g. WO 9515315-318 as described in said patents can be used, too.

- In compounds (V Ac5) the residue of 6-[2-(3-ethyl-2,3-dihydro-thiazolyl)thio-5-methansulphonamido-3H-isobenzofuran-1-one was substituted according to the present invention, has been shown.

This can be prepared according to patent WO 9623786 herein incorporated by reference. Any equivalent pro-

duct containing various substituents as described in said patent can be used, too.

- In compounds (V Ad1) the residue of bumetanide 3-(Aminosulfonyl)-5-(butylamino)-4-phenoxybenzoic acid was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3,806,534 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ad2) the residue of ticrynafen [2,3-Dichloro-4-(2-thienylcarbonyl)-phenoxy]acetic acid was substituted according to the present, has been shown.

This can be prepared according to patent USP 3,758,506 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (VAd3) the residue of ethacrynic acid [2,3-Dichloro-4-(2-methylene-1-oxobutyl)phenoxy]acetic acid was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3,255,241 herein incorporated by reference. Any equivalent pro-

duct containing various substituents as described in said patent can be used, too.

- In compounds (V Ad4) the residue of piretanide 3-(Aminosulfonyl)-4-phenoxy-5-(1-pyrrolidinyl)benzoic acid was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 4,010,273 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae1) the residue of tripamide (3 α , 4 α , 7 α , 7 α)-3-(Aminosulphonyl)-4-chloro-N-octaidro-4,7-methano-2H-isoindol-2-yl)benzamide was substituted according to the present invention, has been shown.

This can be prepared according to patent JP 73 05, 585 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae2) the residue of torsemide N-[[[(1-Methylethyl)amino]carbonyl]4-[(3-methylphenyl)amino]-3-pyrinesulfonamide was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 4,018,929

herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae3) the residue of azosemide 2-Chloro-5-(1H-tetrazol-5-yl)-4-[(2-thienylmethyl)amino]benzenesulphonamide was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3,665,002 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae4) the residue of bendroflumethiazide 3,4-Dihydro-3-(phenyl-methyl)-6-(trifluoromethyl)-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3,392,168 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae5) the residue of chlorothiazide 6-Chloro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide was substituted according to the present inven-

tion, has been shown.

This can be prepared according to patent USP 2,809,194, USP 2,937,169 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae6) the residue of hydrochlorotiazide 6-Chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide was substituted according to the present invention, has been shown.

This can be prepared according to patent DE 1,163,332, USP 3,043,840 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae7) the residue of methyclothiazide (6-Chloro-3-(chloromethyl)-3,4-dihydro-2-methyl-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide was substituted according to the present invention, has been shown.

This can be prepared according to patent Close et al., J.Am.Chem.Society 82, 1132 (1960) herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae8) the residue of chlorthalidone 2-Chloro-5-(2,3-dihydro-1-hydroxy-3-oxo-1H-isoindol-1-yl)benzensulfonamide was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3,055,904 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae9) the residue of Indapamide 3-(Aminosulfonyl)-4-chloro-N-(2,3-dihydro-2-methyl-1H-indol-1-yl)benzamide was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3,565,911 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae10) the residue of metolazone 7-Chloro-1,2,3,4-tetrahydro-2-nethyl-3-(2-methylphenyl)-4-oxo-6-quinazolinesulfonamide was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 3,360,518 herein incorporated by reference. Any equivalent product containing various substituents as described in

said patent can be used, too.

- In compounds (V Ae11) the residue of quinetazone 7-Chloro-2-ethyl-1,2,3,4-tetrahydro-4-oxo-6-quinazolin-sulfonamide was substituted according to the present invention, has been shown.

This can be prepared according to patent USP 2,976,289 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

- In compounds (V Ae12) the residue of furosemide 5-(Aminosulfonyl)-4-chloro-2-[(2-furanylmethyl)amino]benzoic acid was substituted according to the present invention, has been shown.


This can be prepared according to patent USP 3,058,882 herein incorporated by reference. Any equivalent product containing various substituents as described in said patent can be used, too.

The compounds under V Ad/V Ae are particularly suitable for the treatment of urinary disorders, in particular of anuresis.



X₁, in formula A-X₁-NO₂, is a bivalent connecting bridge chosen from the following:

- YO


where Y is a linear or whenever possible branched C_1 - C_{20} alkylene, preferably having from 2 to 5 carbon atoms, or an optionally substituted cycloalkylene having from 5 to 7 carbon atoms;



where n_3 is an integer from 0 to 3;



where nf' is an integer from 1 to 6, preferably from 2 to 4;



wherein R_{1f} = H, CH_3 and nf is an integer from 1 to 6, preferably from 2 to 4.

The processes for obtaining the compounds which contain R from groups I A - IV A are described in patent application WO 95/30641 herein incorporated by reference.

The processes for preparing the compounds of class V A are those described above in application WO 95/30641.

The products of the invention are therapeutically useful in the treatment of various forms of urinary incontinence at lower doses than the corresponding precursor products without the NO donor group and with a wider activity spectrum and without causing the disadvantage previously described for this kind of precursors.

It has been surprisingly found by the Applicant that, meaningfully, the products of the invention do not show reduced pharmacological activity compared to precursors. Conversely, they have a wider pharmacological range of action, since a synergy between the cyclooxygenase inhibiting effect and the myorelaxing effect related to the opening of potassium channels and/or release of nitric oxide, was unexpectedly observed in the lower urinary tract. The products of the invention exhibit a higher safety and do not induce tachyphylaxis.

Additionally, the Applicant found that the products of the invention carry out a pharmaco-therapeutic activity in diverse appropriate experimental models, as described below:

- articular inflammation (musculoskeletal disease) in rats; see Winter C. et al., *Caraggeenin-induced edema*

in hind paw of the rat as an assay for antiinflammatory drugs, Proceedings of the Society for Experimental Biology and Medicine 1962, 111, 544-47;

- *respiratory disease for example bronchospasm from bradykinin in Guinea pigs (Del Soldato P. et al., The anesthetized Guinea pig as a versatile pharmacological test object, Jour. of Pharmacological Methods, 1981 5, 279-285);*
- *vascular disease, such as re-stenosis induced in rats (Role of kinins and nitric oxide in the effects of angiotensin converting enzyme inhibitors on neointima formation, Fahry-RD et al., CIRC-RES. 72/6 (1202-1210)1983);*
- *gynaecological and obstetrical diseases:*
as shown in hyperexcitability states in rat isolated myometrium (Izumi H. et al., Gestational changes in L-arginine-induced relaxation of pregnant rat and nonpregnant myometrial contractility, Am. J. Obstet. Gynecol., 1993, 169, 1327-1337);
- *blood platelet aggregation in women in a pre-eclampsia condition (Janes SI et al., Flow cytometric detection of circulating activated platelets and platelet hyper-responsiveness in pre-eclampsia and pregnancy, Clin.*

Science, 86, 731-739, 1994).


- intestinal tumours, such as for example in experimental adenocarcinoma in rats (Dubois R. et al., *Increased cyclooxygenase-2 levels in carcinogen-induced rat colonic tumors*, Gastroenterology, 110, 1259-1262, 1996).

Therefore, based on the experimental results obtained the above products may be therapeutically useful in the following diseases, in addition to urinary incontinence:


- musculoskeletal disease of an inflammatory nature: group V A;
- respiratory disease, for example bronchitis, in particular asthma, etc.: compounds of the groups from I A to V A;
- gynaecological and obstetrical diseases, including premature delivery, pre-eclampsia and dysmenorrhoea: groups from I A to V A and, additionally, the compounds from group VI A as defined below;
- vascular disease such as re-stenosis: compounds from groups I A to VI A;
- gastrointestinal tumour: compounds from groups from I A to VI A.

The compounds in group VI A, where $t = 1$, include the

following:



(Ia)



(Ib)

where:

R_1 is group $OCOR_3$; where R_3 is methyl, ethyl or a linear or branched C_3 - C_5 alkyl, or the residue of a single-ring heterocycle having 5 or 6 atoms which can be aromatic, partially or totally hydrogenated, containing one or more heteroatoms independently chosen from O, N and S;

R_2 is hydrogen, hydroxy, halogen, a linear or whenever possible branched alkyl having from 1 to 4 C atoms, a linear or whenever possible branched alcoxyl having from 1 to 4 C atoms; a linear or whenever possible branched perfluoroalkyl having from 1 to 4 C atoms, for example trifluoromethyl,

nitro, amino, mono-or di(C₁₋₄)alkylamino;

R₁ and R₂ jointly are the dioxymethylene group, with the proviso that when X = NH, then X₁ is ethylene and R₂ = H; R₁ cannot be OCOR₃ at position 2 when R₃ is methyl; nI being an integer from 0 to 1;

preferably in Ia), X is equal to O or NH, R₁ is acetoxy, preferably at position 3 or 4, most preferably in the ortho position to CO. X₁ is ethylene or (CH₂CH₂O)₂, R₂ is hydrogen or halogen, most preferred are the following A X₁ NO₂ compounds: 3-acetoxy-N-(2-nitroxyethyl)-benzamide, 4-acetoxy-N-(2-nitroxyethyl)-benzamide, 3-acetoxy-N-(5-nitroxypenthyl)-benzamide, 2-acetoxy-N-(5-nitroxypenthyl)-benzamide, N-2-(nitroxyethyl)-2-propionoxybenzamide, 2-acetoxy-2-nitroxyethylbenzoate, 2-acetoxy-N-(cis-2-nitroxycyclohexyl)-benzamide, 2-acetoxy-4-chloro-N-(2-nitroxyethyl)-benzamide, N-(2-nitroxyethyl)-2-((4-thiazolindinyl)carbonyloxy)-benzamide hydrochloride, 2-nicotinoyloxy-N-(2-nitroxyethyl)-benzamide, 2-acetoxy-5-nitroxypenthylbenzoate;

preferably in Ib) R₃ = CH₃, nI = 0;

X is equal to O, X₁ is ethylene; in this case Ib) is the residue of acetylsalicylsalicylic acid.

The processes to obtain the compounds which contain R in group VI A are described in patent WO 95/30641 herein in-

incorporated by reference.

The examples below are intended as an explanation not a limitation of the present invention.

EXAMPLES

Examples 1,2,3 and from 1A to 1F (comparison)

Chemical synthesis

The following compounds were prepared: NO-indomethacin (NO-I), NO-flufenamic (NO-F), NO-nimesulide (NO-M), NO-Naproxen (NO-N).


Preparation of NO-Indomethacin (NO-I)

| | |
|---------------------------------|--------|
| 3-Hydroxybenzyl nitrate | 9.5 g |
| Indomethacin | 7.4 g |
| Dicyclohexylcarbodiimide | 5.6 g |
| CH ₂ Cl ₂ | 200 ml |

were reacted and the solution was allowed to react overnight at room temperature, concentrated to a small volume and filtered. The filtrate was dried and passed through a column containing gel by using chloroform/ethyl acetate 14:1 as an eluting system. A head fraction was thus separated and purified by chromatography using a 2-mm silica plate. Each plate was run three times in a mobile phase made up of cyclohexane/ethyl acetate 6:1.

A yield of 85% was obtained of indomethacin-NO in group

IV A where R is residue (IV) of indomethacin; $t = 1$; $X = O$ and X_1 is the connecting bridging, shown after YO, where $n_3 = 0$, and having the general formula:



Preparation of NO-Flufenamic (NO-F)


| | | |
|---------------------------------|-----|----|
| 3-Hydroxybenzyl nitrate | 6 | g |
| Flufenamic acid | 13 | g |
| Dicyclohexylcarbodiimide | 9.5 | g |
| CH ₂ Cl ₂ | 150 | ml |
| Ethyl ether | 50 | ml |

were reacted and it was allowed to react overnight, concentrated to a small volume and dicyclohexylurea was filtered. The filtrate was dried and passed through a column containing silica by using CH₂Cl₂ as an eluant. A head fraction was thus separated. This fraction was purified by chromatography using a 2-mm silica plate and a cyclohexane/ethyl acetate 6:1 system. Each plate was run three consecutive times. The

head fraction was recovered by extraction with ethyl ether. The ethereal extract was brought to dryness and gives a yellow oil and a yield of 80% for flufenamic-NO.

The ^1H NMR analysis (CDCl_3 , 200 MHz) gave the following data: 5.5 (2H, s); 6.9 (1H, t); 7.4 (10H, m); 8.2 (1H, dd).

The product obtained has the formula:



Preparation of No-Nimesulide (NO-M)

Preparation of the brominated derivative

N-[(2-PHENOXY-4-NITRO) PHENYL] -N- (6-BROMO) HEXANOYLMETHANE-SULPHONAMIDE

4,85 g 6-Bromohexanoylchloride (23 mmol) was added dropwise to a mixture kept at 0°C of 7 g nimesulide (23 mmol) and 6.4 ml triethylamine (46 mmol) in dichloromethane (80 ml). After stirring for one hour at 0°C , a thin layer chromatography analysis (eluant: toluene/ethyl acetate 9:1) showed that unreacted nimesulide was still present. 1 g acyl chloride (4,7 mmol) and 3 ml triethylamine (22 mmol) were

added to the reaction mixture, the temperature was allowed to rise to room temperature and the reaction mixture was stirred overnight. A chromatographic control showed that the reaction was complete. The reaction mixture was treated with water (50 ml), the organic phase was then washed three times with water (50 ml for each washing), then with diluted NaCH (5% w/v), then dried over anhydrous sodium sulphate (10 g). Solvent evaporation at reduced pressure left a yellow solid residue which was ground twice with two portions of ethyl ether (50 ml each). The air-dried solid was 8.3 g, which corresponds to a yield of 74% and exhibited a melting point of 98°C.

Preparation of NO-Nimesulide (NO-M)

N-[(2-PHENOXY-4-NITRO) PHENYL]-N-(6-NITROXY)HEXANOYLMETHANE-SULPHONAMIDE

A solution of 4 g N-[(2-phenoxy-4-nitro)phenyl]-N-(6-bromo)hexanoyl-methanesulphonamide (8.24 mmol) and 2.8 g silver nitrate (16.48 mmol) in anhydrous acetonitrile (20 ml) was reacted with stirring for 2 days. Then 1 g of silver nitrate (6 mmol) was then added and stirring was continued for another day. The precipitate was removed by filtration and the solvent was evaporated from the filtrate at reduced pressure. The residue was dissolved in a mixture of

equal proportions of ethyl acetate and isopropyl ether and stirred for a few minutes with chromatographic-grade silica gel (5 g). The solid was removed by filtration and the filtrate of the solvent was removed at reduced pressure. The residue was a yellow oil which solidified in time (2.6g). The solid was ground with ethyl ether and dried, and exhibited a melting point of 96°C.

The ^1H NMR spectrum (CDCl_3) showed the following signals:

8.05 (1 H, m); 7.62 (2 H, m); 7.48 (2 H, m); 7.32 (1 H, m);
7.08 (2 H, m); 4.40 (2 H, t); 3.40 (3 H, s); 2.24 (2 H, t);
2.18 (3 H, s); 1.70 (4 H, m); 1.45 (2 H, m).

Preparation of compound NO-Naproxen (NO-N)

Compound NO-Naproxen was prepared according to Example 1h (Example 1) in patent WO 95/30641.

Pharmacological tests

The products were administered in a suspension of carboxymethyl cellulose in in-vivo experiments, while they were dissolved in dimethylsulphoxide in in-vitro studies.

The same vehicle used in the corresponding treatment groups was always used for control groups.

The acute toxicity was roughly determined administering an oral dose of 50 mg/kg of substance to groups of 10 mice.

Death rate and appearance of toxic symptoms were evaluated in a period of 14 days from dosing: no toxic effects were observed at the dose administered.

Contraction inhibiting activity in isolated rat detrusor

Male Wistar rats weighing 200 to 300 g were used. The method used is described by Zhou Q. et al. (1995) (see Example 13). After sacrificing the rats by cervical displacement the urinary bladder was isolated and horizontal strips of detrusor muscle about 2 mm wide and about 5 mm long were obtained from the median region. The strips were placed in baths for isolated organs containing Krebs liquid and subjected to a 1 g tension. Tension variations during the test were measured isometrically by using a pressure transducer connected to a polygraph. The inhibitory effect of a pre-treatment with the test derivatives on contraction induced by 40 mM KCl was determined versus drugs having an opening potassium channel activity (cromakalin, nicorandil) nitroderivatives (nitroglycerin, nicorandil) and anti-inflammatories (indomethacin, naproxen, nimesulide). The results are shown in Table 1.

Table 1

| Example | Product | No. of tests | Inhibition% |
|---------------|------------------------------|--------------|-------------|
| 1A comparison | Cromakalim 10_{-6} M | 10 | 33.3 |
| 1B " " | Nitroglycerin 10_{-5} M | 10 | 28.7 |
| 1C " " | Nicorandil 10_{-4} M | 10 | 26.4 |
| 1D " " | Indomethacin 10_{-4} M | 10 | 38.5 |
| 1E " " | Naproxen $5 \cdot 10_{-4}$ M | 10 | 15.2 |
| 1F " " | Nimesulide 10_{-4} M | 10 | 41.8 |
| 1 | NO-I 10_{-4} M | 10 | 46.3 |
| 2 | NO-N $5 \cdot 10_{-4}$ M | 10 | 31.3 |
| 3 | NO-M 10_{-4} M | 10 | 48.1 |

All new nitroderivatives (Examples 1 to 3) proved to be more active than the products used as comparison.

Examples 4-5 and 4A-4C (comparison)

In vivo studies in normal urinary bladder of conscious rats

Cystometrograms of conscious rats were determined according to the method described by Howe B.B et al. (1995) (see Example 9).

Male Wistar rats weighing about 500 g were used. The rats were anaesthetised with Nembutal. After opening their abdomen and exposing their urinary bladder, a catheter filled with physiological solution was implanted in the urinary bladder and caused to emerge from the back of the animals. The abdominal muscle and skin were then sutured. 48 hours

after surgery the animals were placed in metabolic cages and the catheters were connected to a perfusor which perfused 0.18 ml/min of a physiological solution into the urinary bladder, and to a pressure transducer in order to measure intravesical pressure. After stabilisation for 60 minutes, the animals were orally treated with the test products and urination frequency was than measured during 4 hours after dosing. Table 2 shows the results obtained expressed as a ratio versus the baseline frequency recorded before dosing (IC = interval between contractions).

Table 2

| Example | Treatment | No. of tests | IC treated/IC baseline |
|----------------|-------------------------|--------------|------------------------|
| 4 A | Controls | 8 | 1.05 |
| 4 B comparison | Flufenamic acid 5 mg/kg | 8 | 1.42 |
| 4 | NO-F 5 mg/kg | 8 | 1.62 |
| 4 C comparison | Indomethacin 5 mg/kg | 8 | 1.34 |
| 5 | NO-I 5 mg/kg | 8 | 1.48 |

Both new derivatives (Examples 4-5) proved to be more active than the products used as comparison.

Examples 5-6 and 5A-5B (comparison)

In vivo studies in normal urinary bladders of anaesthetised rats

40 Sprague Dawley rats weighing about 300 g were ran-

domly divided into 4 groups and orally treated twice a day for 4 days according to the following experimental scheme:

1. Controls: 0.5% carboxymethyl cellulose
2. Indomethacin 3 mg/kg
3. NO-I 3 mg/kg
4. NO-F 5 mg/kg

18 hours after the last treatment, the effects on the urinary bladder voiding reflex were evaluated using the method described by Maggi C.A. et al., *Prostanoids modulate reflex micturition by acting through capsaicin-sensitive afferents*, European Journal of Pharmacology, 105-112, 1988.

The animals were anaesthetised with urethane, the urinary bladder was prepared for intraluminal pressure measurement. After a stabilisation period with an empty urinary bladder, this was progressively filled with a physiological solution by slow infusion (0.046 ml/min). A contraction of the urinary bladder was observed upon reflex triggering.

The volume of physiological solution and intraluminal pressure required to evoke the reflex (volume and pressure thresholds) were measured. Table 3 shows the pressure and volume threshold values after treatment, calculated considering 100 the values obtained in control animals. All tested products increased this threshold and can, therefore, be

considered useful in case of detrusor instability.

Table 3

| Example | Treatment | No. of animals | Pressure threshold (%) | Volume threshold (%) |
|---------|--------------|----------------|------------------------|----------------------|
| 5 A | Controls | 10 | 100 | 100 |
| 5 B | Indomethacin | 10 | 190 | 198 |
| 5 | NO-1 | 10 | 223 | 226 |
| 6 | NO-F | 10 | 203 | 205 |

Examples 7-8 (7A-7D as comparison)

In vitro studies in unstable urinary bladder

The vesical hypertrophy model secondary to urethral obstruction in rats described by Malmgren A. et al.: *Cystometrical evaluation of bladder instability in rats with intravesical outflow obstruction*, The Journal of Urology, 1987, 137, 1291-1294, was used in order to evaluate the effect of the drugs on hyperactive vesical muscle.

Male Sprague Dawley rats weighing about 250 g were used. In order to obtain partial urethral obstruction, the rats were anaesthetised with Nembutal and the urinary bladder and urethra were exposed by abdominal incision. A ligature was made around the urethra in the presence of an intraluminal cannula with 1 mm diameter. After suturing the abdominal wall the animals were stabulated for 6 weeks in order for vesical hypertrophy to start.

The in vitro experiments were conducted with the parallel use of strips obtained from normal rats and rats with vesical hypertrophy.

The in vitro urinary bladder strips were prepared as described above and the inhibition induced by the drugs on contraction induced by 1/7 Hz electrical stimulation lasting 1 msec., an above maximal voltage, produced by two platinum electrodes, was measured.

The following table shows the percentage of contraction induced by electrical stimulus in normal and hypertrophic urinary bladders in the presence of the test drugs.

Table 4

| Example | Product/Tissue | No. of tests | Contraction % |
|---------|---|--------------|---------------|
| 7 A | Cromakalim 10^{-6} M (normal) | 6 | 50.5 |
| 7 B | Cromakalin 10^{-6} M (hypertrophic) | 6 | 35.7 |
| 7 C | Indomethacin 10^{-6} M (normal) | 6 | 78.2 |
| 7 D | Indomethacin 10^{-6} M (hypertrophic) | 6 | 76.3 |
| 7 | NO-I 10^{-6} M (normal) | 6 | 61.5 |
| 8 | NO-I 10^{-6} M (hypertropic) | 6 | 40.3 |

Differently from indomethacin, the products with an opening of potassium channel activity and the new compounds were found to be more active in inhibiting hypertrophic uri-

nary bladder contraction than normal urinary bladder.

Examples 9-10 and from 9A to 9B (comparison)

In vivo studies in normal urinary bladder of conscious dogs

The cystometrogram of conscious dogs was determined in accordance with the method described by Howe B. B. et al., ZENECA ZD 6169: a NOVEL K_{ATP} Channel opener with in vivo selectivity for urinary bladder, Journal of Pharmacology and Experimental Therapeutics, 274, 884-890, 1995.

Female Beagle dogs with urinary bladder catheterised through the urethra by operating in sterile conditions, were used. Catheters were connected to a perfusor which perfused into the urinary bladder a physiological solution and to a pressure transducer in order to measure intravesical pressure. After 15 minute stabilisation, a 30 ml bolus of physiological solution was perfused into the urinary bladder in order to measure increased intravesical pressure and a series of smaller boluses were then perfused until spontaneous contractions were observed. After a period of contraction stabilisation, contracting activity was monitored for 60 minutes. The animals were then treated orally with the test products and urination frequency was then measured during 4 hours following dosing in control rats and treated rats. Table 5 shows the results obtained expressed as a ratio ver-

sus the baseline frequency recorded before dosing (IC = interval between contractions).

Table 5

| Examples | Treatment | No. of animals | IC treated/IC baseline |
|-----------------|----------------------------|-----------------------|-------------------------------|
| 9 A comparison | Controls | 5 | 1.03 |
| 9 B comparison | Cromakalim 0.5 mg/kg | 5 | 1.48 |
| 9 C comparison | Flufenamic acid 3 mg/kg | 5 | 1.42 |
| 9 | NO-F 3 mg/kg | 5 | 1.76 |
| 9 D comparison | Indomethacin 3 mg/kg | 5 | 1.25 |
| 10 | NO-I 3 mg/kg | 5 | 1.43 |

Examples 11-12 and 11A-11D (comparison)

Relaxing effect in pig urethral smooth muscle

The method described by Werkström et al., *Factors involved in the relaxation of female pig urethra evoked by electrical field stimulation*, British Journal of Pharmacology, 116, 1599-1604, 1995, was used for sample preparation. Samples of urethra were removed from female pigs about 6 months old.

The urethra was opened longitudinally and samples of smooth muscle about 1x2x6 mm in size were removed from an area about 4 mm below ureteral orifices. The samples of smooth muscle were placed in baths for isolated organs, incubated at 37°C and subjected to a 10 mN tension and con-

nected to a force transducer for measuring mechanical activity. After a period of balancing of about 60 minutes, the prepared samples were exposed to Krebs solution without Ca_{++} to determine the highest relaxation level. Normal tone was then restored by adding Krebs solution. The relaxation effects of the test derivatives were then measured. The test was repeated two consecutive times for each prepared sample in order to evaluate any tachyphylaxis effects. The table below shows the relaxation percentages obtained following the two treatments with each test product, expressed considering 100% the highest relaxation determined by the medium without Ca^{++} .

Table 6

| Example | Product | No. of tests | Relaxation % 1 | Relaxation % 2 |
|---------|-----------------------------|--------------|-------------------|-------------------|
| 11 A | Indomethacin 10^{-5} M | 6 | 1.0 | 1.2 |
| 11 | NO-I 10^{-5} M | 6 | 39.3 | 37.2 |
| 11 B | Flufenamic acid 10^{-5} M | 6 | 12.2 | 13.2 |
| 12 | NO-F 10^{-5} M | 6 | 45.8 | 52.1 |
| 11 C | Nitroglycerin 10^{-5} M | 6 | 32.1 | 7.3 |
| 11 D | L-arginine 10^{-5} M | 6 | 22.7 | 12.2 |

The results show that, while drugs with an anti-inflammatory activity such as indomethacin were practically inactive except for flufenamic acid which has itself a myorela-

xing activity, and conventional NO donors, such as nitroglycerin and arginine, were active but induced tachyphylaxis, the new derivatives which are an object of the invention were active and did not induce any tachyphylaxis.

Examples 13-15 and 13A-13B (comparison)

Relaxing activity on vessel smooth muscle

Male Wistar rats weighing 200 to 300 g were used. The method used is described by Zhou Q. et al. *The inhibitory mechanism of nicorandil in isolated rat urinary bladder and femoral artery*, European Journal of Pharmacology, 153-159, 1995. After sacrificing the rats by cervical displacement, the femoral arteries were isolated for the preparation of helical strips about 1x15 mm in size, from which the endothelium was removed. The prepared strips were placed in baths for isolated organs containing Krebs liquid and subjected to a weight of 0.5 g. Tension variations during the test were isometrically measured by means of a pressure transducer connected to a polygraph. The inhibitory effect of a treatment with the test derivatives on contractions induced by 3×10^{-5} M phenylephrine versus reference preparations having a potassium channel opening activity and/or NO donors was measured.

The results are included in Table 7.

Table 7

| Example | Product | No. of tests | Inhibition % |
|-----------------|------------------------------------|--------------|--------------|
| 13 A comparison | Cromakalim 3×10^{-7} M | 10 | 54.1 |
| 13 B comparison | Nicorandil 10^{-6} M | 10 | 32.6 |
| 13 | NO-I 10^{-4} M | 10 | 22.2 |
| 14 | NO-N $5 \cdot 10^{-4}$ M | 10 | 29.0 |
| 15 | NO-M 10^{-4} M | 10 | 19.5 |

All new compounds proved to be less active than Cromakalim and Nicorandil, even used at higher concentrations, than those shown in specific models (see for example Table 6).

Example 16-17 and 16A-16B (comparison)

In vivo gastrointestinal safety studies

Forty Sprague Dawley rats weighing about 300 g were randomly divided into 4 groups and orally treated twice a day for 4 days according to the following experimental scheme:

1. Controls: carboxymethyl cellulose (0.5% by weight):
(5 ml/kg)
2. Indomethacin 3 mg/kg
3. NO-I 3 mg/kg
4. NO-F 5 mg/kg

Eighteen hours after the last treatment the rats were sacrificed to determine any gastrointestinal damage. No gross changes were observed in the gastroenteric tract of control animals.

In the animals treated with indomethacin ulceration was observed in the stomachs and, additionally, the intestines of most animals (7/19) and in some cases (3/10) even diffuse adhesions. In the group treated with NO-I, only gastric ulcers were observed in 1 animal, and in the group treated with NO-F one animal with a gastric ulcer and an animal with a duodenal ulcer were found.

Examples 18-18A and 18-B (comparison)

Studies of nitroxysynthetase activity

The nitroxy-synthetase inhibiting activity induced by lipopolisaccharide (LPS) was determined in rat neutrofiles after administration of any of the test compounds and compared with that obtained after treatment with the suspending vehicle alone (0.5% carboxymethyl cellulose, 5 ml/kg) and a product used as comparison. Briefly, Wistar rats fasted for 24 hours before treatment received one of the test compounds (10 mg/kg) intraperitoneally or the vehicle LPS (5 mg/kg) intravenously (caudal vein).

Four hours later the animals were sacrificed. Blood was

collected for neutrofil isolation.

The enzymatic activity was determined according to the method described by Assreuy J. et al. *Feedback inhibition of nitric oxide sinthase activity by nitric oxide*, British Journal of Pharmacology, 883-837, 1993.

As shown in Table 8, the test product was found to be very effective in inhibiting nitroxy sinthetase compared to the group treated with the vehicle alone and differently from the reference flufenamic.

Table 8

| EXAMPLE | COMPOUND | DOSE (mg/kg/i.p.) | NITROXY-SYNTHETASE ACTIVITY ^a |
|---------|------------|----------------------|--|
| 18 A | Vehicle | --- | 100 |
| 18 B | Flufenamic | 10 | 98 |
| 18 | NO-F | 10 | 63 |

^a percentage compared to the group treated with the vehicle alone.

Conclusions from the whole tests

The derivatives of the invention were found to be active in several tests aimed at determining the potential pharmacological activity controlling urination.

It should also be noted that the derivatives of the invention were also found to be effective in a broader series of tests than that in which each reference drug was found to be active, confirming the hypothesis that these derivatives are endowed with a superior overall pharmacolo-


gical activity in controlling urinary incontinence.

Furthermore, the derivatives of the invention were found to be better tolerated than the reference products. They appeared to be less harmful to the stomach than the corresponding anti-inflammatory agents and less hypotensive than the standard agents with vasorelaxing activity.

The combined characteristics mentioned above make the products of the invention superior to the reference agents.

CLAIMS

1. Use of the following groups of compounds, or their compositions, for the preparation of medicaments for the treatment of urinary incontinence, having the general formula:




or their salts, where:

A = R(COX)_t where t is an integer 0 or 1;


X = O, NH, NR_{1C} where R_{1C} is a linear or branched alkyl having from 1 to 10 C atoms;

R is chosen from the following groups:

* Group I A), where t = 1,



(IAa)



(IAb)

where:

R_{II5} is H, a linear or whenever possible branched C₁-C₃ alkyl;

R_{II6} has the same meanings as R_{II5}, or when R_{II5} is H it can be benzyl;

R_{III1}, R_{III2} and R_{III3} are equal or different one from the other and are hydrogen, linear or whenever possible branched C₁-C₆ alkyl or C₁-C₆ alkoxy, or Cl, F, Br;

R_{III4} is R_{III1} or bromine;


preferred are the compounds where R_{III1}, R_{III2} and R_{III4} are H, and R_{III3} is Cl and R_{III3} is in the ortho position to NH; R_{II5} and R_{II6} are H, X is equal to O, and X₁ is (CH₂-CH₂-O)₂;

(I Ab) is the residue of 2-[[2-methyl-3-(trifluoromethyl)phenyl]amino]-3-pyridinecarboxylic acid and when -COOH is present it is known as flunixin.

The compounds preferred are those where X = O;

* II A) chosen from the following:

where, when t = 1, R is




where R_{2a} and R_{3a} are H, a linear or whenever possible


branched substituted or non-substituted C_1 - C_{12} alkyl, allyl, with the proviso that when one of the two is allyl the other is H; preferably R_{2a} is H, alkyl has from 1 to 4 C atoms, R_{3a} is H;

R_{1a} is chosen from


II Aa)




(II)




(XXI)




(IV)




(VII)




(XXXV)




(VI)




(VIII)



(IX)



(X)



where meanings are as follows:

- in the compounds of formula (IV), residue of ketoprofen:

R_{III1} is H, SR_{III3} where R_{III3} contains from 1 to 4 C linear or whenever possible branched C atoms;

R_{III2} is H, hydroxy;

preferred are the compounds where R_{III1} and R_{III2} are H, R_{3a} is H, and R_{2a} is methyl, $X = O$;

- in the compounds of formula (XXI), residue of carprofen:

R_{xxio} is H, a linear or whenever possible branched alkyl having from 1 to 6 carbon atoms, a C_1 - C_6 alkoxy-carbonyl bound to a C_1 - C_6 alkyl, a C_1 - C_6 carboxyalkyl, a C_1 - C_6 alkanoyl, optionally substituted with halogen, benzyl or halobenzyl, benzoyl or halobenzoyl;

R_{xxi} is H, halogen, hydroxy, CN, a C_1 - C_6 alkyl optionally containing OH groups, a C_1 - C_6 alkoxy, acetyl, benzoyloxy, SR_{xxi2} where R_{xxi2} is a C_1 - C_6 alkyl; a perfluoroalkyl having from 1-3 C atoms, a C_1 - C_6 carboxyalkyl optio-

nally containing OH groups, NO₂, sulphamoyl, dialkyl sulphamoyl with the alkyl having from 1 to 6 C atoms, or difluoroalkylsulphonyl with the alkyl having from 1 to 3 C atoms;

R_{xxi1} is halogen, CN, a C₁-C₆ alkyl containing one or more OH groups, a C₁-C₆ alkoxy, acetyl, acetamido, benzoyloxy,

SR_{III3} is as above defined, a perfluoroalkyl having from 1 to 3 C atoms, hydroxy, a carboxyalkyl having from 1 to 6 C atoms, NO₂, amino, a mono- or dialkylamino having from 1 to 6 C atoms, sulphamoyl, a dialkyl sulphamoyl having from 1 to 6 C atoms, or difluoroalkylsulphamoyl as above defined; or R_{xxi} jointly with R_{xxi1} is an alkylene dioxy having from 1 to 6 C atoms;

preferred are the compounds where R_{xxio} is H, the connecting bridge is at position 2, R_{xxi} is H, R_{xxi1} is chlorine and is in the para position to nitrogen;

R_{3a} is H, R_{2a} is methyl and X is O;

- in the compounds of formula (XXXV), residue of thia-
profenic acid: Ar is phenyl, hydroxyphenyl optionally
mono- or polysubstituted with halogen, an alkanoyl or
alkoxy having from 1 to 6 C atoms, a trialkyl having
from 1-6 C atoms, preferably from 1-3 C atoms, cyclo-

pentyl o-hexyl o-heptyl, heteroaryl, preferably thienyl, furyl optionally containing OH, pyridyl;

the preferred compounds of formula (XXXV) are those where Ar is phenyl, R_{3a} is H, R_{2a} is methyl and X is O;

- in the compounds of formula (II), residue of suprofen,

the preferred, where $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$;

- in the compounds of formula (VI),

of which the preferred, indoprofen, when R_{2a} is CH_3 or indobufen, when R_{2a} is equal to H and $R_{3a} = CH_3$ and $X = O$;

- in the compounds of formula (VIII),

of which the preferred, etodolac, when $R_{3a} = R_{2a} = H$ and $X = O$;

- in the compounds of formula (VII),

of which the preferred, fenoprofen, when $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$;

- in the compounds of formula (III),

of which the preferred, fenbufen, when $R_{3a} = R_{2a} = H$ and $X = O$;


- in the compounds of formula (X), residue of tolmetin,

when $R_{3a} = R_{2a} = H$ and $X = O$;


- in the compounds of formula (IX), residue of flurbi-

profen, when $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$;


II Ab) :




IIIa)




(XXX)




(XXXI)




(XXXII)



(XXXIII)



(XXXVI)




(XXXVII)

where the meanings are as follows:

- when IIIa) contains $-\text{CH}(\text{CH}_3)-\text{COOH}$ it is known as pranoprofen: α -methyl-5H-[1] benzopyran [2,3-b]pyridine-7-acetic acid; preferred $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$ and $\text{X} = \text{O}$;
- when residue (XXX) contains $-\text{CH}(\text{CH}_3)-\text{COOH}$ it is known as bermoprofen: dibenz [b,f] oxepin-2-acetic acid, preferred is $\text{X} = \text{O}$, $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$;




- residue (XXXI) is known as CS-670: 2-[4-(2-oxo-1-cyclohexylidenemethyl)phenyl]propionic acid, when the radical is $-\text{CH}(\text{CH}_3)-\text{COOH}$; preferred $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$ and $\text{X} = \text{O}$;
 - residue (XXXII) derives from the known pemedolac which contains group $-\text{CH}_2\text{COOH}$, preferred $\text{R}_{2a} = \text{R}_{3a} = \text{H}$ and $\text{X} = \text{O}$;
 - when residue (XXXIII) is saturated with $-\text{CH}_2\text{COOH}$ it is known as pyrazolac: 4-(4-chlorophenyl)-1-(4-fluorophenyl)3-pyrazolyl acid derivatives; preferred $\text{R}_{2a} = \text{R}_{3a} = \text{H}$ and $\text{X} = \text{O}$;
 - when residue (XXXVI) is saturated with $-\text{CH}(\text{CH}_3)-\text{COO}-$ it is known as zaltoprofen. When the residue is saturated with a hydroxy or amine group or the acid salts, the compounds are known as dibenzothiepin derivatives. Preferred $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$ and $\text{X} = \text{O}$;
 - when residue (XXXVII) is CH_2-COOH it derives from the known mofezolac: 3,4-di(p-methoxyphenyl)isoxazol-5-acetic acid; preferred are $\text{R}_{2a} = \text{R}_{3a} = \text{H}$, $t = 1$, $\text{X} = \text{O}$.
- * Group IIIA), where $t = 1$,



where:

R_{IVd} and R_{IVd1} are at least one H and the other a linear or whenever possible branched C_1 - C_6 alkyl, preferably C_1 and C_2 , or difluoroalkyl with the alkyl having from 1 to 6 C atoms, preferred is C_1 , or R_{IVd} and R_{IVd1} jointly form a methylene group;

R_{IV} has the following meaning:



where the compounds of group IIIA) have the following meanings:

- in the compounds of formula (II):

R_{IV-II} is an alkyl having from 1 to 6 C atoms, a cycloalkyl having from 3 to 7 C atoms, an alcoxymethyl having from 1 to 7 C atoms, a trifluoroalkyl having from 1 to 3 C atoms, vinyl, ethynyl, halogen, an alkoxy having from 1 to 6 C atoms, a difluoroalkoxy with the alkyl having from 1 to 7 C atoms, an alcoxymethyloxy having from 1 to 7 C atoms, an alkylthiomethyloxy with the alkyl having from 1 to 7 C atoms, an alkylmethylthio with the alkyl having from 1 to 7 C atoms, cyano, difluoromethylthio, a substituted phenyl- or phenylalkyl with the alkyl having from 1 to 8 C atoms; preferably R_{IV-II} is CH_3O , R_{IVd} is H and R_{IVd1} is CH_3 , and is known as the residue of naproxen;


$X = NH$ and X_1 is equal to $(CH_2)_4$ or $(CH_2CH_2O)_2$; also preferred is the same compound where X is equal to O;

- in the preferred compounds of formula (X), for which the residue of loxoprofen has been shown, R_{IVd} is H and R_{IVd1} is CH_3 , $X = NH$ or O and X_1 is equal to $(CH_2)_4$ or $(CH_2CH_2O)_2$;

- in the compounds of formula (III):


R_{IV-III} is a C_2 - C_5 alkyl, even branched when possible, a C_2 and C_3 alkyloxy, allyloxy, phenoxy, phenylthio, a cycloalkyl having from 5 to 7 C atoms, optionally sub-

stituted at position 1 by a C₁-C₂ alkyl;
 preferred is the compound where R_{IV-III} is



and R_{IVd} = H, R_{IVd1} is CH₃, a compound known as the residue of ibuprofen; X = NH and X₁ is equal to (CH₂)₄ or (CH₂CH₂O)₂; also preferred is the same compound where X = O;

* Group IV A)



(IV)


where A = RCOO, t = 1,

of which the residue of the known indomethacin has been shown.


* Group V A) chosen from the following:

- V Aa) fenamates chosen from the following,


where t = 1




(V Aa1)




(V Aa2)




(V Aa3)




(V Aa4)




- V Ab), derivatives of niflumic acid, where $t = 1$:




- V Ac), COX₂ inhibitors, where $t = 0$ and R is as follows:







(V Ac2)






(V Ac3)




(V Ac4)




- V Ad) derivatives of diuretics when $t = 1$ and R is as follows:






(V Ad4)


- V Ae) derivatives of diuretics when $t = 0$ and R is as follows:




(V Ae1)




(V Ae2)




(V Ae3)




(V Ae4)




(V Ae5)




(V Ae6)




(V Ae7)




(V Ae8)




(V Ae9)



(V Ae10)



(V Ae11)



(V Ae12)

where the meaning in group V A) is as follows:

- in compounds (V Aa1) the residue of enfenamic acid, 2-[(2-phenylethyl)amino]benzoic acid, has been shown;
- in compounds (V Aa2) the residue of flufenamic acid, 2-[[3-(trifluoromethyl)phenyl]-amino]benzoic acid, has been shown;
- in compounds (V Aa3) the residue of meclofenamic acid, 2-[(2,6-dichloro-3-methylphenyl)amino]benzoic acid, has been shown;
- in compounds (V Aa4) the residue of mefanamic acid, 2-[(2,3-dimethylphenyl)amino]benzoic acid, has been shown;
- in compounds (V Aa5) the residue of tolfenamic acid, 2-[(3-chloro-2-methylphenyl)amino]benzoic acid, has been shown;
- in compounds (V Ab1) the residue of niflumic acid, 2-[[3-(trifluoromethyl)phenyl]amino]-3-pyridine carboxylic acid, has been shown;
- in compounds (V Ac1)_{R_{vac1}} attached to the oxygen atom in position 2 of the benzene ring of N-(4-nitrophenyl)methansulphonamide can be phenyl or cyclohexane. When R_{vac1} is phenyl the residue is that of nimesulide;
- in compounds (V Ac2) the residue of 3-formylamino-7-

methylsulfonylamino-6-phenoxy-4H-1-benzopyran-4-one has been shown;

- in compounds (V Ac3) the atom X_4 that links the radical 2,4-difluorothiophenyl to position 6 of the indanone ring of the residue 5-methanesulfonamido-1-indanone can be sulfur or oxygen;

- in compounds (V Ac4) the residue of celecoxib 4-[5-(4-methylphenyl)-3-(trifluoromethyl)pyrazol-1-yl] benzenesulphonamide, has been shown;

- in compounds (V Ac5) the residue of 6-[2-(3-ethyl-2,3-dihydro-thiazolyl)thio-5-methanesulphonamido-3H-isobenzofuran-1-one has been shown.

- in compounds (V Ad1) the residue of bumetanide 3-(Aminosulfonyl)-5-(butylamino)-4-phenoxybenzoic acid has been shown;

- in compounds (V Ad2) the residue of ticrynafen [2,3-Dichloro-4-(2-thienylcarbonyl)-phenoxy]acetic acid has been shown;

- in compounds (V Ad3) the residue of ethacrynic acid [2,3-Dichloro-4-(2-methylene-1-oxobutyl)phenoxy]acetic acid, has been shown;

- in compounds (V Ad4) the residue of piretanide 3-(Aminosulfonyl)-4-phenoxy-5-(1-pyrrolidinyl)benzoic

acid has been shown.

- in compounds (V Ae1) the residue of tripamide (3 α , 4 α , 7 α , 7a α)-3-(Aminosulphonyl)-4-chloro-N-(octahydro-4,7-metano-2H-isoindol-2-yl)benzamide has been shown.
- in compounds (V Ae2) the residue of torsemide N-[[[(1-Methylethyl)amino]carbonyl]4-[(3-methylphenyl)amino]-3-pyridinesulfonamide has been shown;
- in compounds (V Ae3) the residue of azosemide 2-Chloro-5-(1H-tetrazol-5-yl)-4-[(2-thienylmethyl)amino]benzenesulphonamide has been shown;
- in compounds (V Ae4) the residue of bendroflumethiazide 3,4-Dihydro-3-(phenyl-methyl)-6-(trifluoromethyl)-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has been shown;
- in compounds (V Ae5) the residue of chlorothiazide 6-Chloro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has been shown;
- in compounds (V Ae6) the residue of hydrochlorothiazide 6-Chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has been shown;
- in compounds (V Ae7) the residue of methylclothiazide (6-Chloro-3-(chloromethyl)-3,4-dihydro-2-methyl-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has

been shown;

- in compounds (V Ae8) the residue of chlorthalidone 2-Chloro-5-(2,3-dihydro-1-hydroxy-3-oxo-1H-isoindol-1-yl)benzensulfonamide has been shown;

- in compounds (V Ae9) the residue of Indapamide 3-(Aminosulfonyl)-4-chloro-N-(2,3-dihydro-2-methyl-1H-indol-1-yl)benzamide has been shown;

- in compounds (VAe10) the residue of metolazone 7-Chloro-1,2,3,4-tetrahydro-2-methyl-3-(2-methylphenyl)-4-oxo-6-quinazolinesulfonamide has been shown;


- in compounds (V Ae11) the residue of quinetazone 7-Chloro-2-ethyl-1,2,3,4-tetrahydro-4-oxo-6-quinazoline-sulfonamide has been shown;

- in compounds (V Ae12) the residue of furosemide 5-(Aminosulfonyl)-4-chloro-2-[(2-furanylmethyl)amino]benzoic acid has been shown.



X_1 in formula A- X_1 -NO₂ is a bivalent connecting bridge chosen from the following:

- YO


where Y is a linear or whenever possible branched C₁-C₂₀ alkylene, preferably having from 2 to 5 carbon atoms, or an optionally substituted cycloalkylene having from 5 to 7 carbon atoms;



where n_3 is an integer from 0 to 3;



where nf' is an integer from 1 to 6, preferably from 2 to 4;



where $\text{R}_{1f} = \text{H}, \text{CH}_3$ and nf is an integer from 1 to 6, preferably from 2 to 4.

2. Use of the compounds according to Claim 1, in which R is chosen from groups IV A) and V A).
3. Compounds or their compositions for use as medicaments from group V A) in Claim 1.
4. Compounds from group V A) according to Claim 1.
5. Compounds or their compositions for use as medicaments

from group V A) according to Claim 3 for the treatment of musculoskeletal disease of an inflammatory nature, respiratory disease of an inflammatory nature, gynaecological and obstetrical disease including early delivery, pre-eclampsia and dysmenorrhoea, cardiovascular disease including re-stenosis, gastrointestinal tumours.

6. Use of the following compounds, or their compositions, for the preparation of medicaments for the following therapeutical applications:


treatment of respiratory disease: bronchitis, in particular asthma: groups from I A) to V A) in Claim 1;

gynaecological and obstetrical disease including early delivery, pre-eclampsia and dysmenorrhoea: groups from I A) to V A) in Claim 1 and group VI A) as defined below;


vascular disease including re-stenosis: groups from I A) to V A) in Claim 1 and group VI A);

gastrointestinal tumours: groups from I A) to V A) in Claim 1 and group VI A);


the compounds in group VI A) have the general formula



of Claim 1, where $t = 1$, include the following:



(Ia)



(Ib)

where:

R_1 is group $OCOR_3$; where R_3 is methyl, ethyl or a linear or branched C_3 - C_5 alkyl, or the residue of a single-ring heterocycle having 5 or 6 atoms which can be aromatic, partially or totally hydrogenated, containing one or more heteroatoms independently chosen from O, N and S;
 R_2 is hydrogen, hydroxy, halogen, a linear or whenever possible branched alkyl having from 1 to 4 C atoms, a linear or whenever possible branched alcoxyl having from 1 to 4 C atoms; a linear or whenever possible branched perfluoroalkyl having from 1 to 4 C atoms, for example trifluoromethyl, nitro, amino, mono- or

di(C₁₋₄)alkylamino;

R₁ and R₂ jointly are the dioxymethylene group, with the proviso that when X = NH, then X₁ is ethylene and R₂ = H; R₁ cannot be OCOR₃ at position 2 when R₃ is methyl; nI being an integer from 0 to 1;

preferably in Ia), X is equal to O or NH, R₁ is acetoxy, preferably at position 3 or 4, most preferably in the ortho position to CO. X₁ is ethylene or (CH₂CH₂O)₂, R₂ is hydrogen or halogen, most preferred are the following A X₁ NO₂ compounds: 3-acetoxy-N-(2-nitroxyethyl)-benzamide, 4-acetoxy-N-(2-nitroxyethyl)-benzamide, 3-acetoxy-N-(5-nitroxypentyl)-benzamide, 2-acetoxy-N-(5-nitroxypentyl)-benzamide, N-2-(nitroxyethyl)-2-propionoxybenzamide, 2-acetoxy-2-nitroxyethylbenzoate, 2-acetoxy-N-(cis-2-nitroxcyclohexyl)-benzamide, 2-acetoxy-4-chloro-N-(2-nitroxyethyl)-benzamide, N-(2-nitroxyethyl)-2-((4-thiazolindinyl)carbon-yloxy)-benzamide hydrochloride, 2-nicotinoyloxy-N-(2-nitroxyethyl)-benzamide, 2-acetoxy-5-nitroxypentylbenzoate;


preferably in Ib) R₃ = CH₃, nI = 0;

X is equal to O, X₁ is ethylene; in this case Ib) is the residue of acetylsalicylsalicylic acid.



maryd

\\server\name

PSCRIPT Page Separator

[Main Indexes](#)   

[owl home](#) || [writing lab and owl](#) || [handouts](#) || [workshops and presentations](#) || [internet resources](#)

 [Get a printer-friendly version of this page.](#)
 [Get an Adobe PDF version of this page.](#)

A or An? Exercises



Brought to you by the [Purdue University Online Writing Lab](#)

In the following phrases, supply either *a* or *an*:

1. ____ bingo game
2. ____ idiot
3. ____ good job
4. ____ rotten plum
5. ____ used fork
6. ____ uncle
7. ____ historian
8. ____ apple
9. ____ hair
10. ____ artichoke
11. ____ horrible movie
12. ____ opera
13. ____ fine opera
14. ____ television
15. ____ earthquake
16. ____ icicle
17. ____ plant
18. ____ eggplant

19. ____ honorable discharge
20. ____ intelligent man
21. ____ table
22. ____ up stairway
23. ____ paper clip
24. ____ animal
25. ____ usual feeling
26. ____ interest
27. ____ alibi
28. ____ early bird
29. ____ couch
30. ____ airplane
31. ____ grade
32. ____ pair
33. ____ idea
34. ____ energy level

Check your answers!

 [Get a printer-friendly version of this page.](#)
 [Get an Adobe PDF version of this page.](#)

[owl home](#) || [writing lab and owl](#) || [handouts](#) || [workshops and presentations](#) || [internet resources](#)

This page is located at <http://owl.english.purdue.edu/handouts/esl/esliartEX1.html>



Receive weekly tips and updates from Purdue's OWL and Writing Lab!

Copyright ©1995-2004 by OWL at Purdue University and Purdue University. All rights reserved.

Use of this site constitutes acceptance of our [terms and conditions of fair use](#).

To contact OWL, please refer to our [contact information page](#) so that you email the appropriate person. Requests for help sent to the wrong address may not receive a timely response.

maryd

\\server\name

PSCRIPT Page Separator



[owl home](#) || [writing lab and owl](#) || [handouts](#) || [workshops and presentations](#) || [internet resources](#)

Contents

Grammar

**Adjectives
and adverbs
Nouns
Prepositions**

**Pronouns
Sentence
structure
Verbs**

Punctuation

**Apostrophes
and
Quotation
Marks
Commas
Hyphens
Other
punctuation
Sentence
Punctuation**

Spelling

**Sound-alike
words
Noun
Plurals
Suffixes**

Exercises

**Grammar
Spelling
Punctuation**

Grammar, Punctuation, and Spelling

Brought to you by the [Purdue University Online Writing Lab](#).

In this section of our site, we offer you handouts and exercises on grammar, spelling, and punctuation. We also have [PowerPoint presentations](#) related to grammar, and we have an entire section of handouts and resources for [English as a Second Language](#) learners that might also prove useful.

We now have printer friendly versions and Adobe PDF versions of all of these handouts available. Visit our [printer-friendly grammar, spelling, and punctuation index](#) to download and print any of these handouts, or click on the link on any of the handouts to visit the printer-friendly version.

Grammar

Adjectives and adverbs

- [A or An?](#): with exercise and answer key
- [Adjective or Adverb?](#): with 2 exercises and answer keys
- [Adjectives with Countable and Uncountable Nouns](#)
- [Using Articles \(A/An/The\)](#): with 2 exercises and answer keys

Nouns

- [A Little Help with Capitals](#)
- [Count and NonCount Nouns \(with Plurals, Articles, and Quantity Words\)](#): with 2 exercises and answer keys
- [Count and NonCount Nouns \(with Articles and Adjectives\)](#): with exercise and answer key
- [Writing Numbers](#): with exercise and answer key

Prepositions

- [Prepositions of Direction: To, On\(to\), In\(to\)](#) with exercise and answer key
- [Prepositions of Location: At, In, On](#) with exercise and answer key
- [Prepositions of Time, of Place, and to Introduce Objects](#)
- [Spatial Relationships of Prepositions](#)

Pronouns

- [Using Pronouns Clearly](#)
- [Pronoun Case](#)

Sentence structure

- [Appositives](#) : with exercise and answer key
- [Dangling Modifiers](#)
- [Independent and Dependent Clauses \(definitions\)](#)
- [Making Subject and Verbs Agree](#)
- [Parallel Structure](#)
- [Run-ons, Comma splices, and Fused Sentences](#)
- [Sentence Fragments](#): with 3 exercises and answer keys
- [Sentence Punctuation Patterns](#)
- [Some Strategies for Improving Sentence Clarity](#)

Verbs

- [Active/Passive Verbs](#)
- [Irregular Verbs](#)
- [Sequence of Tenses](#)
- [Tense Consistency](#): with an exercise and answer key
- [Two-Part \(Phrasal\) Verbs- \(Idioms\)](#)
- [Verb Tenses](#)
- [Verbs with Helpers \(Perfectives and Auxiliaries\)](#)
- [Verbals: Gerunds, Participles, and Infinitives](#)
- [Voice and Mood](#)

[Return to Top](#)

Punctuation

Apostrophes and Quotation Marks

- [Apostrophes](#) with exercise and answer key
- [Quotation Marks](#): with exercise and answer key

Commas

- [Commas after Introductions](#): with exercise and answer key
- [Commas vs. Semicolons in Compound Sentences](#) :
with exercise and answer key
- [Commas with Essential and Non-essential Elements](#): with 3
exercises and answer keys
- [Proofreading for Commas](#)

- [Using Commas](#):with 5 exercises and answer keys

Hyphens

- [Using Hyphens](#)

Other punctuation

- [Brief Overview of Punctuation: Semicolons, Colons, Parentheses, Dash, Quotation Marks, and Italics](#): with exercise and answer key

Sentence Punctuation

- [Sentence Punctuation Patterns](#)
- [Run-ons, Comma splices, and Fused Sentences](#)

[Top of the page](#)

Spelling

- [Spelling: Sound-Alike Words](#)
- [Spelling: *Accept/Except* and *Affect/Effect*](#): with 2 exercises and answer keys
- [Spelling: i/e rules](#): with 2 exercises and answer keys
- [Spelling: Noun Plurals](#)
- [Spelling: Suffixes--*ible* vs. *able*](#): with exercise and answer key

[top of the page](#)

Exercises

Try our [interactive exercises index](#) for interactive versions of some of these exercises.

- [A or An?](#)
- [Adjective or Adverb?](#)
 - [Exercise 1](#)
 - [Exercise 2](#)
- [Apostrophes](#)
- [Appositives](#)
- [Articles \(A/An/The\)](#):
 - [Exercise 1](#)
 - [Exercise 2](#)
- [Commas](#):
 - [Exercise 1](#)
 - [Exercise 2](#)

- [Exercise 3](#)
 - [Exercise 4](#)
 - [Exercise 5](#)
- [Commas after Introductions](#)
- [Commas Set Off Nonessential Elements:](#)
 - [Exercise 1](#)
 - [Exercise 2](#)
 - [Exercise 3](#)
- [Commas vs. Semicolons in Compound Sentences](#)
- [Exercises for Eliminating Wordiness](#)
- [Count and NonCount Nouns Exercises](#)
- [Quantity Terms with Count and Noncount Nouns](#)
- [Writing Numbers:Exercise](#)
- [Practice Exercises in Paraphrasing](#)
- [Making Subjects and Verbs Agree](#)
- [Prepositions of Direction:To, On \(to\), In \(to\)](#)
- [Prepositions of Location: At, In, On](#)
- [Punctuation Exercise 1](#)
- [Quotation Marks](#)
- [Sentence Fragments](#)
 - [Exercise 1](#)
 - [Exercise 2](#)
 - [Exercise 3](#)
- [Spelling: accept/except](#)
 - [Exercise 1](#)
 - [Exercise 2](#)
- [Spelling: i/e rules](#)
 - [Exercise 1](#)
 - [Exercise 2](#)
- [Spelling:Exercises for -ible vs. —able](#)
- [Tense Consistency](#)

[Top of the page](#)

[owl home](#) || [writing lab and owl](#) || [handouts](#) || [workshops and presentations](#) || [internet resources](#)

This page is located at <http://owl.english.purdue.edu/handouts/index.html>



Receive weekly tips and updates from Purdue's OWL and Writing Lab!




Copyright ©1995-2004 by OWL at Purdue University and Purdue University. All rights reserved.
Use of this site constitutes acceptance of our [terms and conditions of fair use](#).

To contact OWL, please refer to our [contact information page](#) so that you email the appropriate

person. Requests for help sent to the wrong address may not receive a timely response.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|----|---|
| <p>(51) International Patent Classification ⁶ : C07C 203/04, 211/46, 311/00, C07D 209/08, 209/48, 209/72, 213/00, 213/74, 231/12, 239/91, 285/28, 295/14, 307/52, 311/00, 311/22, 333/22, 409/12, 417/12, A61K 31/215, 31/245, 31/405, 31/63</p> | A3 | <p>(11) International Publication Number: WO 98/09948</p> <p>(43) International Publication Date: 12 March 1998 (12.03.98)</p> |
| <p>(21) International Application Number: PCT/EP97/04774</p> <p>(22) International Filing Date: 2 September 1997 (02.09.97)</p> <p>(30) Priority Data: MI96A001821 4 September 1996 (04.09.96) IT</p> <p>(71) Applicant (for all designated States except US): NICOX S.A. [FR/FR]; 45, avenue Kléber, F-75116 Paris (FR).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): DEL SOLDATO, Piero [IT/IT]; Via Toti, 22, I-20052 Monza (IT). SANNICOLÒ, Francesco [IT/IT]; Alzaia Naviglio Grande, 46, I-20148 Milano (IT).</p> <p>(74) Agents: SAMA, Daniele et al.; Sama Patents, Via G.B. Morgani 2, I-20129 Milano (IT).</p> | | <p>(81) Designated States: AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p> <p>(88) Date of publication of the international search report: 4 June 1998 (04.06.98)</p> |
| <p>(54) Title: NITRIC ESTER DERIVATIVES AND THEIR USE IN URINARY INCONTINENCE AND OTHER DISEASES</p> <p>(57) Abstract</p> <p>Use of the following groups of compounds or their compositions for the preparation of medicaments for the treatment of urinary incontinence, such compounds having general formula: A-X₁-NO₂ or their salts, where A = R(COX)_t and where t is an integer 0 or 1; X = O, NH, NR_{1C}, where R_{1C} is a linear or branched alkyl having from 1 to 10 C atoms; R is (IA) where t = 1 and X₁ is equal to -YO- where Y is a C₁-C₂₀ alkylene, C₅-C₇ cycloalkyl or oxyalkyl derivatives.</p> <div style="text-align: right;">  <p>(IA)</p> </div> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/04774

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C203/04 C07C211/46 C07C311/00 C07D209/08 C07D209/48
 C07D209/72 C07D213/00 C07D213/74 C07D231/12 C07D239/91
 C07D285/28 C07D295/14 C07D307/52 C07D311/00 C07D311/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|---|-----------------------|
| X | ROUSSEAU P ET AL: "URINARY INCONTINENCE IN THE AGED PART 2: MANAGEMENT STRATEGIES" GERIATRICS, vol. 47, no. 6, January 1992, pages 37-40,47, XP002050834 * p.47, 2nd col., Functional incontinence, 1st par. * | 1,2 |
| X | WO 95 30641 A (NICOX LTD; DEL SOLDATO PIERO (IT); SANNICOLA FRANCESCO (IT)) 16 November 1995 cited in the application * p.5, last par.-p.6, l.12; p.53, last par-p.58, bottom; claims 1-7 * --- -/-- | 1-6 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

24 March 1998

Date of mailing of the international search report

09. 04. 98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Uiber, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/04774

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D333/22 C07D409/12 C07D417/12 A61K31/215 A61K31/245
A61K31/405 A61K31/63

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | WO 94 04484 A (CORLAY SL ;METGROVE LTD (IE); MATJI JOSE ANTONIO (ES); ALCAIDE ANT) 3 March 1994 * p.2, 1.12-20; claims 1-7 * | 5,6 |
| Y | see claims 1-7 | 1-4 |
| A | WO 94 21618 A (CERMOL SA ;SUNKEL CARLOS (ES); FAU MIGUEL (ES); PRIEGO JAIME G (ES) 29 September 1994 see claims 1-10 | 1,2 |
| A | WO 92 01668 A (ITALFARMACO SPA) 6 February 1992 see the whole document | 1,2 |
| P,Y | WO 97 25984 A (SCHERING AG ;UNIV TEXAS (US)) 24 July 1997 * p.4, 1.21-29; claims 1 and 8 * | 1,2 |
| | -/- | |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

24 March 1998

Date of mailing of the international search report

09.04.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Uiber, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/04774

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|---|-----------------------|
| X | US 5 439 938 A (SNYDER SOLOMON H ET AL) 8 August 1995 see claims 1,24 --- | 1,2,5,6 |
| X | US 5 480 999 A (CHABRIER DE LASSAUNIERE PIERRE ET AL) 2 January 1996 * col.1, 1.11-col.2, 1.43; claims 1-14 * --- | 3-6 |
| X | WO 94 13635 A (MERCK FROSST CANADA INC ;FORD HUTCHINSON ANTHONY W (CA); KENNEDY B) 23 June 1994 cited in the application * p.13, 1.20-p.14, 1.6; p.26, compound 12 * --- | 3-6 |
| P,X | WO 97 16405 A (NICOX S A ;DEL SOLDATO PIERO (IT); SANNICOLA FRANCESCO (IT)) 9 May 1997 see claims 1-6 --- | 6 |
| Y | WO 95 13802 A (SCHERING AG ;GARFIELD ROBERT E (US); YALLAMPALLI CHANDRA (US)) 26 May 1995 * p.5, 1.16; claims 1-3 * --- | 3-6 |
| Y | WO 96 16645 A (WELLCOME FOUND ;KING S COLLEGE LONDON (GB); BELDER ADAM JULIAN DE) 6 June 1996 * p.1, 1st and 2nd par.; p.3, 1.10-13; claims 1-4 * --- | 3-6 |
| Y | WO 96 17604 A (WELLCOME FOUND ;KING S COLLEGE LONDON (GB); BELDER ADAM JULIAN DE) 13 June 1996 * p.1, 1st par.; p.3, 1.10-11; claims 1-20 * --- | 3-6 |
| Y | K.F. CHUNG : "Furosemide and other diuretics in asthma" J. ASTHMA, vol. 31, no. 2, 1994, pages 85-92, XP002059907 * Table 1; p.90, conclusion * --- | 3-6 |
| Y | CIRINO ET AL: "Inhibition of inducible NO synthetase expression by novel NSAID derivatives with gastrointestinal sparing properties" BRITISH J. PHARMACOLOGY, vol. 117, 29 March 1996, pages 421-26, XP002059908 see abstract --- -/-- | 1-6 |

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/EP 97/04774

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|---|-----------------------|
| Y | WO 96 24585 A (SEARLE & CO ;LEE LEN F (US)) 15 August 1996 * p.4, 1.7-13, 1.18-19; p.29, 1.12-15; claims 1-4 * | 3-6 |
| Y | ANDERSSON K -E ET AL: "NITRIC OXIDE SYNTHASE AND THE LOWER URINARY TRACT: POSSIBLE IMPLICATIONS FOR PHYSIOLOGY AND PATHOPHYSIOLOGY" SCANDINAVIAN JOURNAL OF UROLOGY AND NEPHROLOGY, no. SUPPL. 175, 1995, pages 43-53, XP000670055 see abstract | 1-4 |
| Y | ARA G ET AL: "CYCLOOXYGENASE AND LIPOXYGENASE INHIBITORS IN CANCER THERAPY" PROSTAGLANDINS LEUKOTRIENES AND ESSENTIAL FATTY ACIDS, vol. 54, 1996, pages 3-16, XP002053643 * p.6, A Colon tumors * | 3-6 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 97/04774

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1(partially)-6(partially)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

In view of the great number of compounds for which protection is being sought in claim 1, the search has been restricted for economical reasons to compounds of the examples and the first inventive concept.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/ EP 97 /04774

FURTHER INFORMATION CONTINUED FROM PCT/ISA/

1. Claims: 1,2

Use of the compounds A-X1-NO2, A being R(COX)t and R being choosen from groups I A) to V A) for the preparation of medicaments for treating urinary incontinence

The subject matter of the present application is not unitary contrary to the requirements of R.13. 1 PCT.

With respect to the present claims 1 to 6, the ISA is unable to see any common inventive concept linking these claims as:

The subject matter of claim 1 is directed to the USE of compounds of a general formula A-X1-NO2, with A being R(COX)- and R is choosen from groups I a) to V A) for the manufacture of a medicament for treating URINARY INCONTINENCE.

Claims 3 and 4 are directed to a sub-class of said compounds (or compositions containing them) of the general formula above, R is choosen from group V A)

The subject matter of claim 5 is directed to a sub-class of compounds of a general formula A-X1-NO2, with A being R(COX)- and R is choosen from groups V A)(or compositions containing said compounds) for use as medicament for treating:

- a) musculoskeletal disease of an inflammatory nature
- b) respiratory diseases
- c) gynaecological or obstetrical diseases
- d) vascular diseases
- e) gastrointestinal tumors

The subject matter of claim 6 is directed to the use of compounds of a general formula A-X1-NO2, with A being R(COX)- and R is choosen from groups I a) to VI A) for the manufacture of a medicament for treating diseases b) to e) listed above

With respect to the various definitions of R given in the present application, it appears that they shall necessarily comprise a -O-NO2 moiety as SOLE structural requirement. Organic nitrates comprising the same moiety -O-NO2 are well known in therapy of cardiovascular diseases (see W09201688 or W09421618). The compounds when R belongs to the groups I A) to IV A) and VI A) are already known for therapy (see W09530641).

=> In other words, the present application covers the use of compounds comprising said -O-NO2 moiety for various therapeutic uses.

As some of the compounds are already known in therapy, the ISA comes to the conclusion that the present application is directed to the subsequent therapeutical uses of partially known compounds. The ISA is unable to identify any common inventive concept which links the subsequent claimed

INTERNATIONAL SEARCH REPORT

International Application No. PCT/ EP 97/04774

FURTHER INFORMATION CONTINUED FROM PCT/ISA/

diseases: urinary incontinence and the diseases a)-e) listed above.
Accordingly, the present application is not unitary.

For instance, claim 6 when it refers to the compounds when R belongs to the groups I A) to IV A) and VI A) comprises the following 4 different inventions:

- 1) use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) for the preparation of medicaments for treating respiratory diseases
- 2) use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) for the preparation of medicaments for treating gynaecological or obstetrical diseases
- 3) use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) for the preparation of medicaments for treating vascular diseases
- 4) use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) for the preparation of medicaments for treating gastrointestinal tumors

A comprehensive search for the subsequent inventions would have caused major additional searching efforts

Accordingly, the present application comprises the following 6 inventions:

1st invention:
see above Subject 1

2nd invention
claims 3-5 and 6(partially)
use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups V A) for the preparation of medicaments for treating respiratory diseases, gynaecological or obstetrical diseases, vascular diseases or gastrointestinal tumors

3rd invention:
claim 6(part.)
use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) and VI A) for the preparation of medicaments for treating respiratory diseases

4th invention:
claim 6 (part.)
use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) and VI A) for the preparation of medicaments for treating gynaecological or obstetrical diseases

5th invention:
claim 6(part.):
use of compounds of formula A-X1-NO2 with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) and VI A) for the preparation of medicaments for treating vascular diseases

6th invention:

FURTHER INFORMATION CONTINUED FROM PCT/ISA/

claim 6 (part.)

use of compounds of formula A-X1-NO₂ with A= R(COX)t, R is a moiety according to any of groups I a)- IV A) and VI A) for the preparation of medicaments for treating gastrointestinal tumors

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/04774

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9530641 A | 16-11-95 | IT 1269735 B | 15-04-97 |
| | | IT 1274609 B | 18-07-97 |
| | | AU 2215695 A | 29-11-95 |
| | | AU 678063 B | 15-05-97 |
| | | AU 7809294 A | 01-05-95 |
| | | BR 9407749 A | 12-02-97 |
| | | CA 2173582 A | 13-04-95 |
| | | CA 2190087 A | 16-11-95 |
| | | WO 9509831 A | 13-04-95 |
| | | EP 0722434 A | 24-07-96 |
| | | EP 0759899 A | 05-03-97 |
| | | HU 74446 A | 30-12-96 |
| | | HU 75961 A | 28-05-97 |
| | | JP 9503214 T | 31-03-97 |
| | | JP 9512798 T | 22-12-97 |
| | | US 5700947 A | 23-12-97 |
| WO 9404484 A | 03-03-94 | IT 1256345 B | 01-12-95 |
| | | AT 143941 T | 15-10-96 |
| | | CA 2120942 A | 03-03-94 |
| | | DE 69305322 D | 14-11-96 |
| | | DE 69305322 T | 20-02-97 |
| | | EP 0609415 A | 10-08-94 |
| | | ES 2093979 T | 01-01-97 |
| | | JP 7500355 T | 12-01-95 |
| | | US 5597847 A | 28-01-97 |
| WO 9421618 A | 29-09-94 | CH 685700 A | 15-09-95 |
| | | AU 6153694 A | 11-10-94 |
| | | CA 2135921 A | 29-09-94 |
| | | EP 0640078 A | 01-03-95 |
| | | JP 7506844 T | 27-07-95 |
| WO 9201668 A | 06-02-92 | IT 1243367 B | 10-06-94 |
| | | AT 118478 T | 15-03-95 |
| | | AU 8097491 A | 18-02-92 |
| | | CA 2087442 A | 27-01-92 |
| | | DE 69107459 D | 23-03-95 |
| | | EP 0540544 A | 12-05-93 |
| | | ES 2056783 T | 16-10-94 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/04774

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|--|--|
| WO 9201668 A | | HU 213405 B US 5589490 A US 5366992 A | 30-06-97 31-12-96 22-11-94 |
| WO 9725984 A | 24-07-97 | AU 1703197 A | 11-08-97 |
| US 5439938 A | 08-08-95 | NONE | |
| US 5480999 A | 02-01-96 | AT 401054 B AT 256092 A AU 664399 B AU 3049892 A BE 1006227 A CA 2085555 A CH 685629 A DE 4244539 A DK 157592 A ES 2052452 A FR 2685869 A FR 2685916 A GB 2263111 A,B GR 1001443 B HK 22296 A IE 71675 B IT 1256761 B JP 5286916 A LU 88208 A NL 9300001 A NZ 245499 A PL 169432 B PT 101165 A SE 9203825 A US 5360925 A ZA 9210080 A | 25-06-96 15-10-95 16-11-95 08-07-93 14-06-94 05-07-93 31-08-95 08-07-93 05-07-93 01-07-94 09-07-93 09-07-93 14-07-93 30-12-93 16-02-96 26-02-97 15-12-95 02-11-93 15-04-93 02-08-93 26-07-95 31-07-96 28-02-94 05-07-93 01-11-94 02-08-93 |
| WO 9413635 A | 23-06-94 | US 5604260 A AU 5621594 A CA 2151235 A EP 0673366 A JP 8504408 T | 18-02-97 04-07-94 23-06-94 27-09-95 14-05-96 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/04774

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9716405 A | 09-05-97 | IT M1952263 A | 30-04-97 |
| | | AU 7495096 A | 22-05-97 |
| ----- | | | |
| WO 9513802 A | 26-05-95 | AU 8144594 A | 06-06-95 |
| | | BR 9408067 A | 24-12-96 |
| | | CA 2176721 A | 26-05-95 |
| | | CN 1135178 A | 06-11-96 |
| | | CZ 9601390 A | 16-10-96 |
| | | EP 0730448 A | 11-09-96 |
| | | FI 962109 A | 15-07-96 |
| | | HU 74465 A | 30-12-96 |
| | | JP 9505068 T | 20-05-97 |
| | | NO 961993 A | 12-07-96 |
| | | PL 314465 A | 16-09-96 |
| | | SK 63396 A | 05-03-97 |
| ----- | | | |
| WO 9616645 A | 06-06-96 | AU 3878895 A | 19-06-96 |
| ----- | | | |
| WO 9617604 A | 13-06-96 | AU 3991095 A | 26-06-96 |
| ----- | | | |
| WO 9624585 A | 15-08-96 | US 5596008 A | 21-01-97 |
| | | AU 4770796 A | 27-08-96 |
| | | EP 0808305 A | 26-11-97 |
| ----- | | | |

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.